

Interacting electrons with spin in a one-dimensional dirty wire connected to leads

I. Safi and H.J. Schulz

Laboratoire de Physique des Solides, Université Paris-Sud, 91405 Orsay, France

We investigate a one-dimensional wire of interacting electrons connected to semi-infinite leads in the absence and in the presence of a backscattering potential. An electron incident on the clean wire is perfectly transmitted into spatially separated spin and charge parts in the noninteracting leads, a result we extend to any finite-range interactions. The backscattering potential is renormalized in a non-universal way and therefore the reduction in the conductance is more complicated than the laws derived up to now: it has power laws as a function of temperature, wire length, and also the distance of a barrier to the contacts.

72.10.-d, 73.40.Jn, 74.80.Fp

Recently intensive theoretical and experimental effort has been devoted to the understanding of the transport properties of a Luttinger liquid. The theoretical prediction of conductance renormalization by the interactions has not been observed in quantum wires [1,2]. The physics of such tiny structures has shown that one has to take into consideration the experimental setup. In that context, a model [3,4] simulating the two-dimensional Fermi gas into which a quantum wire opens by perfect one-dimensional leads was proposed; this amounts to retain only the propagating mode. In that geometry, the conductance is not renormalized by the interactions as commonly accepted. The effect of a backscattering potential on the conductance of spinless electrons was discussed in ref. [3,5,6] taking into account contact effects which had been ignored in other work [7,8]. In this letter, we consider the nontrivial effects of the external leads, the contacts, and of spin. We will study the transmission process of an incident spin-polarized electron through the clean wire, demonstrating an interesting consequence of the spin-charge separation typical of the Luttinger liquid. We then treat the effect of weak impurities, showing that their effect on the conductance depends crucially on contact effects, especially when isolated barriers are considered. This may be important in many experimental cases as currently studied wires are often quite short. We give an improved general framework to treat weak disorder at finite temperature and in an inhomogeneous system.

An important low-energy property of an interacting one-dimensional system is the separation of spin and charge dynamics [9–11]. The Hamiltonian is decoupled as $H = H_\rho + H_\sigma$, where:

$$H_\nu = \int \frac{dx}{2\pi} \left[u_\nu K_\nu \Pi_\nu^2 + \frac{u_\nu}{K_\nu} (\partial_x \Phi_\nu)^2 \right] \quad (1)$$

for $\nu = \rho, \sigma$. The boson fields are related to the charge and spin density (ρ and σ) through: $\sqrt{2}\partial_x \Phi_\rho(x)/\pi = \rho$ and $\sqrt{2}\partial_x \Phi_\sigma/\pi = \sigma$. Π_ν is the momentum density conjugate field to Φ_ν . In the absence of interactions, $K_\nu = 1, u_\nu = v_F$. Consider now an interacting wire delimited by $[-a, a]$ connected perfectly to noninteracting leads. The global system thus formed is described by the Hamiltonian (1) with spatially varying parameters $u_\nu(x)$, and $K_\nu(x)$. We set $u_\nu = v_F, K_\nu = 1$ on the external leads, i.e. for $|x| > a$. For simplicity, we consider the case $u(x) = u_\nu$ and $K(x) = K_\nu$ constants on the interacting wire.

An interesting issue we consider first is the transmission process of an incident electron: let us imagine that we inject an electron of definite spin in the left lead, and place a spin and charge detector at some location on the right lead. This corresponds to create a kink in both Φ_ρ and Φ_σ whose time evolution has now to be solved given the initial conditions. The equations of motion for these fields are decoupled, and require their continuity at the contacts as well as that of $u_\nu \partial_x \Phi_\nu / K_\nu$. In particular, both the charge and spin current, $j_\nu = \frac{\sqrt{2}}{\pi} \partial_t \Phi_\nu$ are conserved. This is because we restrict ourselves to interactions which conserve these currents: we suppose that both umklapp processes and as the backscattering of electrons with opposite spin are irrelevant, as is the case for purely repulsive interactions. [9–11]

The injected electron propagates freely in the left lead until it reaches the first contact with the interacting wire. In view of the above continuity requirements, its charge and spin are reflected with two different coefficients γ_ν due to the change in interactions: $\gamma_\nu = (1 - K_\nu)/(1 + K_\nu)$. The transmitted charge and spin propagate at different velocities $u_\rho \neq u_\sigma$, thus reach the second contact where they get partially transmitted at different times: $t_\nu = 2a/u_\nu$. Since the transmitted charge and spin propagate at the same Fermi velocity in the right non-interacting lead, they will stay *spatially separated*, at a distance $v_F |t_\rho - t_\sigma|$ (Fig.1). The remaining charge (spin) in the wire will continue to bounce back and forth at the contacts, letting at each time cycle t_ρ (t_σ) a partial charge (spin) come out. Thus we end up with a series of spatially separated partial charge and spin spikes on the two non interacting leads: they don't recombine as one might expect. The transmitted spikes rather correspond to a complicated superposition of electron-hole excitations. Note that in the relevant case of spin-invariant interactions, $K_\sigma = 1$, the spin part is not reflected but gets directly transmitted to the right lead, while the charge undergoes

the multiple reflection process. At times very long compared to t_ρ (t_σ), the series of transmitted charge (spin) spikes sum up to unity. Thus the transmission of an incident spin polarized flux is perfect. We can generalize the perfect transmission result to any profile of the interaction parameters, and even more, to the case of *Coulomb interactions screened on the measuring leads* [12]. By arguments analogous to those of the spinless case [3,13] the perfect transmission leads to a conductance $g = 2e^2/h$, independent of interactions.

We will now show that in the presence of backscattering by impurities or at the contacts the conductance depends on the interactions, but is generally affected by the external leads. Consider any backscattering potential $V(x)$. The most general representation of the backscattering Hamiltonian δH respecting the initial symmetries is: [14]

$$\delta H \sim \int_{-a}^{+a} dx \sum_{m_\rho, m_\sigma}^l V(x; m_\rho, m_\sigma) e^{2im_\rho \phi_0(x)} e^{i\sqrt{2}(m_\rho \Phi_\rho + m_\sigma \Phi_\sigma)} \quad (2)$$

The sum runs over integers m_ρ and m_σ of the same parity. The function ϕ_0 includes $k_F x$ as well as the effects of the inhomogeneity of the interactions and the forward scattering [12].

For definiteness we specialize to the case of a barrier localized at a point x . In order to derive the renormalization equations, we expand the exact partition function Z at finite temperature in terms of V . Z describes a neutral gas of integer charges restricted to two circles, one corresponding to the charge ($\nu = \rho$) and the other to the spin ($\nu = \sigma$) degree of freedom. The radius of each circle ν is given by β . Two charges on different circles don't interact, but have to be of the same parity if located at the same angle. Two charges on the same circle ν interact via $U_\nu(x, x, \tau) = \langle \Phi_\nu(x, 0) [\Phi_\nu(x, 0) - \Phi_\nu(x, \tau)] \rangle$. U_ν is not the Coulomb interaction, but an infinite series of Logarithmic terms which are related to the transmission process we exposed earlier.

We increase the cutoff to $\tau = \tau_0 e^l$, where τ_0 is the bare cutoff of the order of the inverse of the Fermi energy, and modify the parameters in order to keep Z invariant. The leading-order flow equations at finite temperature are:

$$\frac{dV}{dl} = \left[1 - \frac{1}{2} \left(m_\rho^2 \frac{dU_\rho}{dl} + m_\sigma^2 \frac{dU_\sigma}{dl} \right) \right] V \quad (3)$$

where for simplicity we have omitted the arguments. In particular, through U , V now acquires a dependence on the position of the barrier.

We can show that new interaction terms between the charges are generated by the renormalization, but they are decaying faster than U_ν at long time scales [12]. Thus the equation (3) is sufficient, and can be integrated straightforwardly. In the absence of an external energy

scale, the unique limitation on increasing the cutoff comes naturally from the fact that one has to put at least two charges on the same circle of radius β , thus we cannot go further than $\tau = \beta/2$. We write

$$V^2(x; m_\rho, m_\sigma; l) = V^2(x; m_\rho, m_\sigma) \prod_{\nu=\rho, \sigma} E_\nu(x; m_\rho, m_\sigma; l)$$

where $V(x; m_\rho, m_\sigma; l)$ are the renormalized parameters. It would be cumbersome to give the behavior of E_ν at arbitrary temperatures. We restrict ourselves to the two limits of high or low temperature compared to T_L : here, T_L is a common notation for both temperature scales $1/t_\nu = u_\nu/2a$ for $\nu = \rho, \sigma$. In the limit $T \gg T_L$, the multiple reflections caused by the change in interactions don't affect the correlation functions due to the lack of thermal coherence along the wire. Only one reflection is felt within a thermal length $L_T = v_F/T$ from the contacts. For $\nu = \rho$ or σ , we denote by $t_{x\nu} = (a - |x|)/u_\nu$ the time it takes for a charge or spin excitation emanating at x to reach the closest contact. Up to a slowly varying function, we get:

$$E_\nu \sim (T\tau_0)^{m_\nu^2 K_\nu - 1} \left[(T\tau_0)^2 + \tanh^2 \pi T t_{x\nu} \right]^{m_\nu^2 K_\nu \gamma_\nu / 2} \quad (4)$$

At points far from the contacts compared to L_T , this expression coincides with that obtained in an infinite Luttinger liquid with parameters K_ν :

$$E_\nu \sim (T\tau_0)^{m_\nu^2 K_\nu - 1} \quad (5)$$

At the contacts, a similar expression holds substituting $K_\nu \rightarrow K_{a\nu} = 1 - \gamma_\nu$, which is the charge/spin transmission coefficient through one contact. In the intermediate region $\tau_0 \ll t_{x\nu} \ll \beta$, the expression (4) has an additional power of $t_{x\nu}$.

In the limit of low temperature compared to T_L , the multiple reflections affect the correlation functions, thus the external leads determine the temperature dependence. But there is a nontrivial dependence on the wire length as well as on the distance of the barrier from the contacts:

$$E_\nu \sim (T\tau_0)^{m_\nu^2 - 1} (T_L \tau_0)^{-m_\nu^2 \gamma_\nu} \left[1 + (t_{x\nu}/\tau_0)^2 \right]^{m_\nu^2 K_\nu \gamma_\nu / 2} \quad (6)$$

For a barrier in the bulk, so that $t_{x\nu} \sim t_\nu/2 = a/u_\nu$, this expression simplifies to:

$$E_\nu \sim (T\tau_0)^{m_\nu^2 - 1} (T_L \tau_0)^{m_\nu^2 (K_\nu - 1)} \quad (7)$$

At the contacts, one has to replace K_ν by $K_{a\nu}$. For $t_{\nu x} \gg \tau_0$, the expression (6) contains a power law in $t_{\nu x}$.

We now discuss the effect of the backscattering potential on the conductance. For this purpose, we use the equation of motion of Φ_ρ to derive a Dyson equation for the nonlocal conductivity and thus cast the exact expression for the conductance into a form appropriate for a perturbative computation [5,12]. The latter

is equivalent to inserting the renormalized parameters $V^2(x; m_\rho, m_\sigma; l)$ in the expression for the conductance because the interactions are not renormalized. The dominant correction is reported in table I. It is given by $V^2(x, 1, 1)$, i.e. by the $2k_F$ backscattering, as long as $K_\rho > 1/3$. Consider now the case $K_\rho < 1/3$. In an infinite Luttinger liquid the tendency towards a $4k_F$ CDW dominates over the $2k_F$ CDW, which is reminiscent of a tendency towards a Wigner crystal. Consequently, in this parameter region the $4k_F$ scattering process ($m_\rho = 2$, $m_\sigma = 0$), corresponding to a coherent backscattering of two electrons, becomes important. It turns out that the measuring leads affect this tendency. The dominant correction to the conductance is not necessarily given by the $4k_F$ backscattering term as in an infinite Luttinger liquid. In particular, below T_L the noninteracting leads dominate, saturating the $4k_F$ scattering. Details of the crossover between dominant $2k_F$ and $4k_F$ scattering depend on the temperature as well as on the location, as shown in table I. In particular, the crossover temperature is position-dependent. For $1/5 < K_\rho < 1/3$, this crossover takes place on the bulk but not near the contacts. Thus, the external leads *suppress the importance of $4k_F$ scattering at enough low temperature or close to the contacts* and also saturate the tendency towards the formation of a Wigner crystal.

All the previous results concerning one barrier, including the renormalization equations (3), can be extended to any potential V with a spatial extension much less than the thermal length L_T and the wire length L [12]. Note that in the presence of a potential with any extension, the partition function describes a gas of integer charges confined to two cylinders. The renormalization equations (3) are still valid. Nevertheless, the potential V can renormalize the interactions. The computation of the conductance can be accomplished in a perturbative way [12] as long as the potential is either weak enough or not too much extended to not renormalize the interactions. Now we perform the average over random impurities extended all over the wire. For simplicity, we limit ourselves to the case of a Gaussian distribution: $\langle V(x)V(y) \rangle = v_F^2 \delta(x - y)/l_e$. We perform the average of the conductance expression. Upon integration over the wire, the reduction δg contains an infinity of powers in the temperature. We skip the details, and give the leading correction in table II. There appear essentially two terms in δg : the contribution of the impurities close to the contacts, governed by the local parameter $K_{a\rho}$, and that of the impurities on the bulk governed by K_ρ . The latter dominates the former at any temperature if $K_\rho < 1 + \sqrt{2}$ or at $T > T_b$ if $K_\rho > 1 + \sqrt{2}$, with

$$T_b \tau_0 = (T_L \tau_0)^{-1/(\gamma_\rho K_\rho)}. \quad (8)$$

For $K_\rho < 1/3$, the $4k_F$ backscattering term dominates only at high temperature $T > T_L$.

Let us discuss the conductance fluctuations. We can evaluate the variance of the conductance, and show that:

$$\frac{1}{2}(\delta g)^2 \leq \langle g^2 \rangle - \langle g \rangle^2 \leq (\delta g)^2 \quad (9)$$

As long as δg is small, the conductance is self-averaging, but its fluctuations are of the same order as its reduction.

Let us compare finally the conductance we get to the usual conductance of an infinite dirty Luttinger liquid. While the ballistic conductance we found is not renormalized by the interactions, its reduction in the presence of impurities is similar concerning its power law behavior controlled by the interactions, but is more complicated than this in ref. [15,14,16]. In particular, we must comment on the commonly used scaling argument used to infer the properties of the finite wire from those computed in an infinite wire: The energy cutoff introduced by the temperature T is replaced by the higher one introduced by the wire length T_L . In our model, we observe that the substitution $T \rightarrow T_L$ does not allow to go from (4) to (6) unless $m_\rho = m_\sigma = 1$ and $t_{x\nu} \sim t_\nu$. Thus even in the presence of a barrier at a distance from the contact of the order of the wire length, we recover the correction to the conductance of an infinite wire (extrapolated on the whole range of temperature) only for $K_\rho > 1/3$ (see table I). In the presence of extended disorder, table II shows that the infinite wire result is recovered merely in the window $1/3 < K_\rho < 1 + \sqrt{2}$.

We come now to the comparison with experiment. Recently, Tarucha *et al.* studied relatively clean and long (up to $10\mu\text{m}$, corresponding to $T_L = 0.1\text{K}$) quantum wires fabricated in an AlGaAs/GaAs heterostructure. At high enough temperature, the conductance is quantized in units of $2e^2/h$. As the temperature is lowered, the impurities become more efficient, and the quantization less perfect. The reduction of the one-channel wire conductance with the temperature is attributed to electron-electron interactions and fitted with a power law controlled by a parameter $K_\rho = 0.7$. These experimental observations contradict the result of renormalization of the ballistic conductance by the interactions. Instead, they can be explained in a natural way by the model we proposed. Let's discuss further the backscattering potential. It can emanate either from impurities or from imperfections at the opening of the wire in the two-dimensional gas. Our model yields a peculiar dependence of the reduction to the conductance on the temperature and the location of the barriers. For instance, if the backscattering is due merely to the contacts, the parameter Tarucha *et al.* infer would correspond to the local parameter $K_{a\rho} = 2K_\rho/(1 + K_\rho)$, which yields $K_\rho = 0.5$. On the other hand, they don't have a good agreement with the intuitive law $[T^2 + T_L^2]^{(K_\rho-1)/2}$ [16]. We predict instead different possible functions of the temperature. For instance, in the presence of impurities in the intermediate region, we get two successive power laws controlled first

by K_ρ then by $K_{a\rho}$ upon cooling (table I). We are more inclined to discard the possibility of a Gaussian distribution extended over the wires studied by Tarucha and al. First, the fluctuations in the conductance are not of the same order of magnitude as its reduction, eq.(9). Secondly, the reduction depends on the gate voltage, and shows even some oscillations for the $10\mu m$ wire. Probably, the backscattering potential configuration depends on each wire, and more precise fit and evaluation of K_ρ are required. In particular, one hopes to infer a parameter K_ρ depending on the filling factor.

Another interesting experiment was performed more recently by Yacoby *et al.* [2]. The conductance is again not renormalized by the interactions at high enough temperature. Upon cooling, the conductance decreases, the reduction being more important in the longer wires. This is qualitatively coherent with the predictions we get in the presence of backscattering for $K_\rho < 1$. Nonetheless, the observed reduction in the conductance is independent on the gate voltage, while both K_ρ and the overall scale of δg (which depends on the Fermi energy) are expected to depend on the gate voltage. As discussed in Ref. [2], it's not granted that the leads have a non-Fermi behavior in such sophisticated experimental set up, and a more suited model is needed.

To conclude, we studied the properties of an interacting wire connected to measuring leads in the absence and in the presence of a back scattering potential. The clean wire separates an incident electron into spin and charge parts, and this separation persists even in the noninteracting leads. The total transmission is perfect for any range of the interactions. In the presence of a backscattering potential, the conductance is affected both by the interactions and by the external leads. It's only for certain impurity distributions and interaction parameters that the latter don't intervene.

- [9] V. J. Emery, in *Highly Conducting One-Dimensional Solids*, edited by J. T. Devreese, R. P. Evrard, and V. E. van Doren (Plenum, New York, 1979), p. 247.
- [10] H. J. Schulz, in *Proceedings of Les Houches Summer School LXI*, edited by E. Akkermans, G. Montambaux, J. Pichard, and J. Z.-J. stin (Elsevier, Amsterdam, 1995).
- [11] J. Voit, Rep. Prog. Phys. **58**, 977 (1995).
- [12] I. Safi-Taktak, Ph.D. thesis, Laboratoire de Physique des Solides, 1996.
- [13] R. Landauer, Phil. Mag. **21**, 863 (1970).
- [14] C. L. Kane and M. P. A. Fisher, Phys. Rev. B **46**, 1220 (1992).
- [15] W. Apel and T. M. Rice, Phys. Rev. B **26**, 7063 (1982).
- [16] M. Ogata and H. Fukuyama, Phys. Rev. Lett. **73**, 468 (1994).

FIG. 1. Dynamic transmission of an incident electron with spin up. The charge and spin are separated even in the non-interacting leads. As an example, we consider here $u_\rho > u_\sigma$ and $K_\sigma < 1$, $K_\rho > 1$.

FIG. 2. The reduction to the conductance scaled by the backscattering strength as a function of the temperature for different potential distributions. We choose $K_\rho = 0.5$, $K_\sigma = 1$, and $E_F = 148T_L$, thus $\ln(T_L/E_F) = 5$. The dotted, dashed and dot-dashed curves correspond respectively to a barrier at the center of the wire, at the contact and at an intermediate point x . There is a crossover at T_L from a power law controlled by K_ρ ($K_{a\rho}$) at the center (at the contact) to a plateau. At x , there are two crossovers: one at $T_x = u_\rho/t_x$ from $T^{K_\rho-1}$ to $T^{K_{a\rho}-1}$, then a saturation occurs at T_L . The continuous curve correspond to an extended Gaussian distribution, where the power law at $T < T_L$ is governed by K_ρ since the interactions are repulsive. Since $K_\rho = 0.5 > 1/3$, the $2k_F$ backscattering dominates at all T .

TABLE I. A weak barrier at x : dominant reduction of the conductance depending on the temperature and on K_ρ for the case $K_\sigma = 1$. $T_x = \pi/t_x$ is the inverse of the time to get from x to the closest contact. Every temperature has to be multiplied by τ_0 . The first line gives the behavior all over the wire at $T < T_L$, governed only by the $2k_F$ backscattering. Note that the second line has to be skipped in case $T_x \sim T_L$. The last line concerns only $t_x \gg \tau_0$.

	$K_\rho < 1/5$	$1/5 < K_\rho < 1/3$	$1/3 < K_\rho$
$T < T_L$	$T_L^{-\gamma_\rho} T_x^{-\gamma_\rho K_\rho}$		
$T_L < T < T_x$	$T^{2-4\gamma_\rho} T_x^{-4K_\rho\gamma_\rho}$	$T^{-\gamma_\rho} T_x^{-\gamma_\rho K_\rho}$	
$T_x < T$	$T^{4K_\rho-2}$		$T^{K_\rho-1}$

-
- [1] S.Tarucha, T. Honda, and T. Saku, Sol. State Comm. **94**, 413 (1995).
 - [2] A. Yacoby *et al.*, Phys. Rev. Lett. **77**, 4612 (1996).
 - [3] I. Safi and H. J. Schulz, Phys. Rev. B **52**, R17040 (1995).
 - [4] D. Maslov and M. Stone, Phys. Rev. B **52**, R5539 (1995).
 - [5] I. Safi and H. J. Schulz, *Transport through a single-band wire connected to measuring leads* (Kluwer Academic Press, AA Dordrecht, 1995).
 - [6] I. Safi and H. J. Schulz, in *Correlated Fermions and Transport in Mesoscopic Systems*, edited by T. Martin, G. Montambaux, and J. T. T. Van (Editions Frontieres, Gif-sur-Yvette, 1996).
 - [7] D. Maslov, Phys. Rev. B **52**, R14368 (1995).
 - [8] A. Furusaki and N. Nagaosa, Phys. Rev. B **54**, R5239 (1996).

TABLE II. Gaussian disorder: dominant reduction of the conductance δg multiplied by l_e . The $4k_F$ backscattering dominates for $K_\rho < 1/3$ only at high temperature $T > T_L$. The temperature T_b is given by eq. (8).

	$K_\rho < 1/3$	$1/3 < K_\rho < 1 + \sqrt{2}$	$1 + \sqrt{2} < K_\rho$
$T < T_L$		$T_L^{K_\rho-2}$	$T_L^{-\gamma_\rho}$
$T_L < T < T_b$	$T^{4K_\rho-2}/T_L$	$T^{K_\rho-1}/T_L$	$T^{-\gamma_\rho}$
$T_b < T$	$T^{4K_\rho-2}/T_L$	$T^{K_\rho-1}/T_L$	

